



# New Phases of $\text{QCD}_3$ and $\text{QCD}_4$

Nathan Seiberg

IAS

D. Gaiotto, A. Kapustin, Z. Komargodski, and NS, arXiv:1703.00501;

Z. Komargodski, and NS, arXiv:170?.?????;


D. Gaiotto, Z. Komargodski, and NS, arXiv:170?.?????

# 4d pure gauge $SU(N)$

- Large  $N$ : 1<sup>st</sup> order transition at  $\theta = \pi$ .
  - $CP$  (or  $T$ ) is spontaneously broken there [Witten]
- Finite  $N$ : a one-form global symmetry associated with the center of the gauge group [Gaiotto, Kapustin, NS, and Willett].
  - At  $\theta = \pi$ : mixed 't Hooft anomaly between it and  $CP$ .
  - 't Hooft anomaly matching: cannot move continuously from confinement at  $\theta = 0$  to confinement at  $\theta = 2\pi$ .
  - Specifically, at  $\theta = \pi$  deconfinement, or broken  $CP$ , or TQFT
  - Simplest scenario (as at large  $N$ ): a single 1<sup>st</sup> order phase transition at  $\theta = \pi$ .
  - Assume it (more exotic scenarios are in the paper).

# 4d pure gauge $SU(N)$ at finite $T$

- Because of the anomaly, cannot move continuously from confinement at  $\theta = 0$  to confinement at  $\theta = 2\pi$ .
- Hence,  $T_{deconfinement} \leq T_{CP}$ .



# QCD<sub>4</sub> with one quark ( $N_f = 1$ )

- No chiral symmetry for massless quarks, but at infinite  $N$  a massless  $\eta'$   $m_{\eta'}^2 = \frac{1}{N} \Lambda^2 + O\left(\frac{1}{N^2}\right)$  [Witten]


- With massive quarks

$$m_{\eta'}^2 = \frac{1}{N} \Lambda^2 + \text{Re}(m) \Lambda + O(1/N^2, m^2, m/N)$$

- Therefore, even at finite  $N$  can find an exactly massless  $\eta'$ .
- For large  $|m|$  the same behavior as in the pure gauge system.
- First order transition in the complex  $m e^{i\theta}$  plane

$m e^{i\theta}$

massless  $\eta'$   
“4d Ising point”



# QCD<sub>4</sub> with $N_f$ quarks for $1 < N_f < N_{CFT}$

- For  $N_f \geq N_{CFT}(N)$  a CFT or lack of asymptotic freedom
- For  $1 < N_f < N_{CFT}$  a massless theory –  $SU(N_f)$  sigma model
- Turn on equal masses. As for  $N_f = 1$ , a first order transition line at  $\theta = \pi$ , which ends at the massless point [Dashen]

$$\left| m^{N_f} e^{i\theta} \right|$$



$SU(N_f)$  sigma model

# QCD<sub>3</sub> $SU(N)_k$ with $N_f$ quarks

- Now we can add a Chern-Simons term with coefficient  $k$ .
- For large  $N_f$  a non-trivial fixed point [Appelquist and Nash]
- Assume that this remains the case for  $N_f \geq N^*(N, k)$
- Topological phases at large  $|m|$

$$SU(N)_{k-N_f/2}$$

$$SU(N)_{k+N_f/2}$$


$$m < 0$$



Phase transition, CFT

$$m > 0$$

- What happens for  $N_f < N^*(N, k)$ ?
- Use recently suggested dualities...



# Dual descriptions

Many references. These are some of the recent ones.

...; Giombi, Yin; Aharony, Gur-Ari, Yacoby; Giombi, Minwalla, Prakash, Trivedi, Wadia, Yin; Maldacena, Zhiboedov; Aharony, Giombi, Gur-Ari, Maldacena, Yacoby; Jain, Minwalla, Sharma, Takimi, Wadia, Yokoyama; Minwalla, Yokoyama; Yokoyama; Jain, Mandlik, Minwalla, Takimi, Wadia, Yokoyama; Inbasekar, Jain, Mazumdar, Minwalla, Umesh, Yokoyama; Jain, Minwalla, Yokoyama; Gur-Ari, Yacoby; Son; Wang, Senthil; Metlitski, Vishwanath; Barkeshli, McGreevy; Radicevic; **Aharony**; Karch, Tong; NS, Senthil, Wang, Witten; Hsin, NS; Kachru, Mulligan, Torroba, Wang; Metlitski, Vishwanath, Xu; Aharony, Benini, Hsin, NS; Benini, Hsin, NS ...



# Dual descriptions

$$N_f \text{ fermions coupled to } SU(N)_k \quad \leftrightarrow \quad N_f \text{ scalars at } |\Phi|^4 \text{ point coupled to } U\left(\frac{N_f}{2} + k\right)_{-N}$$

We take  $N$  positive and  $k$  non-negative.

The scalars are in a generalized Wilson-Fisher fixed point or a gauged version of it.

For  $N_f > 2k$  apply time reversal and then  $k \rightarrow -k$  to find another possible duality

$$SU(N)_k \quad \leftrightarrow \quad U\left(\frac{N_f}{2} - k\right)_N$$

# Dual descriptions

$N_f$  fermions coupled to

$N_f$  scalars at  $|\Phi|^4$  point coupled to

- $SU(N)_k \leftrightarrow U\left(\frac{N_f}{2} + k\right)_{-N}$
- $SU(N)_k \leftrightarrow U\left(\frac{N_f}{2} - k\right)_N$

Checks:

- For large  $N$  and  $k$  with fixed ratio explicit calculations and relation to AdS duals (only the first one)
- For  $N_f = 0$  level/rank duality (only the first one)
- Relation to SUSY dualities of [...; Giveon, Kutasov; ...]
- Flow to fewer flavors
- Global symmetry and 't Hooft anomaly matching (for simplicity, limit to  $N > 2$ )

# Problem for $N_f > 2k$

$N_f$  fermions coupled to

$N_f$  scalars at  $|\Phi|^4$  point coupled to


- $SU(N)_k \leftrightarrow U\left(\frac{N_f}{2} + k\right)_{-N}$
- $SU(N)_k \leftrightarrow U\left(\frac{N_f}{2} - k\right)_N$

A mass deformation in the fermionic theory leads to a gapped system  $SU(N)_{k \pm N_f/2}$  (depending on the sign of the mass).


In the scalar theories for one sign it leads to a gapped theory  $U\left(\frac{N_f}{2} + k\right)_{-N} \leftrightarrow SU(N)_{k + N_f/2}$ , which is good.

But with the other sign the gauge symmetry is completely Higgsed and we end up with a massless theory.

$SU(N)_k$  with  $N_f$  quarks  $N_f \leq 2k$



$SU(N)_k$  with  $N_f$  quarks  
 $2k < N_f < N^*(N, k)$



# $SU(N)_k$ with $N_f$ quarks

$$2k < N_f < N^* (N, k)$$

- Three phases
  - For large  $|m|$  semiclassical physics – gapped, topological.
  - For small  $|m|$  a new quantum phase with global symmetry breaking  $U(N_f)/U\left(\frac{N_f}{2} + k\right) \times U\left(\frac{N_f}{2} - k\right)$
  - Each phase transition has a weakly coupled bosonic dual description
- The intermediate phase
  - Wess-Zumino term from the Chern-Simons term
  - For  $k = 0$ :  $U(N_f) \rightarrow U(N_f/2) \times U(N_f/2)$  with a WZ term
  - Assuming this we can derive for other values of  $k$
  - Skyrmions in the nonlinear model are monopoles in the bosonic theory and are the baryons in the fermionic theory

# 4d pure gauge $SU(N)$

Return to 4d. Will soon relate to the 3d story.

Study the domain wall at the first order transition point at  $\theta = \pi$ .

- The theory on the domain wall needs to account for the different 't Hooft anomalies between the two sides of the wall.
- $SU(N)_{-1}$  (Same as on the domain wall between neighboring vacua of  $\mathcal{N} = 1$  SUSY  $SU(N)$  pure gauge theory.)

# QCD<sub>4</sub> with $N_f = 1$

$$me^{i\theta}$$

massless  $\eta'$   
“4d Ising point”

Study the domain wall at the transition.

- No anomaly argument
- For large negative  $me^{i\theta}$ , expect  $SU(N)_{-1}$
- For small mass, should be trivial – use the  $\eta'$  theory
- There must be a phase transition on the domain wall.
- Same phases as in 3d

$$SU(N)_{-1/2} \text{ with } N_f = 1 \quad \psi \leftrightarrow \quad U(1)_N \text{ with } N_f = 1 \quad \phi$$



# QCD<sub>4</sub> with $1 < N_f < N_{CFT}$

$$\boxed{m^{N_f} e^{i\theta}}$$

$SU(N_f)$  chiral  
Lagrangian

Study the domain wall at the transition.

- For large negative  $m^{N_f} e^{i\theta}$  expect  $SU(N)_{-1}$
- For small mass,  $CP^{N_f-1}$  with  $N\Gamma_{WZ}$  – use the chiral Lagrangian
- There must be a phase transition on the domain wall
- Same phases as in 3d

$$SU(N)_{-1+N_f/2} \text{ with } N_f \text{ quarks} \leftrightarrow U(1)_N \text{ with } N_f \text{ scalars}$$

- Consistent with the intermediate phase in the 3d discussion

# Summary

QCD<sub>4</sub>

- $N_f = 0$ 
  - New parity anomaly
  - Phase transition at  $\theta = \pi$  for all  $N$
  - $T_{deconfinement} \leq T_{CP}$
- $N_f = 1$ 
  - massless  $\eta'$  for all  $N$
- All  $N_f$ 
  - first order transition with domain walls

# Summary

QCD<sub>3</sub> with a Chern-Simons term

- Large  $N_f$ : a second order transition separating two gapped topological phases
- Small  $N_f$ : same as large  $N_f$ , but with a bosonic dual
- Intermediate  $N_f$ : three phases. Two of them are gapped and topological. Intermediate phase with global symmetry breaking.

Consistent with the analysis of domain walls in QCD<sub>4</sub>

Interesting generalization to  $SO(N)/Spin(N)$  and  $Sp(N)$  gauge theories – new insights about confinement.